HYDROBIOLOGIA
Volumes 422/423, 1 April 2000

Theme
Assessing the Ecological Integrity of Running Waters

Guest Editors
M. Jungwirth, S. Muhar & S. Schmutz

Symposium sponsors ix

List of authors xi–xii

List of reviewers xiii–xvi

Preamble
M. Jungwirth, S. Muhar, S. Schmutz xvii–xviii

Preface: The Austrian approach
W. Stalzer, H. Blöch xix–xx

Introduction: Sustaining living rivers
J.R. Karr, E.W. Chu 1–14

Basic Components of Running Waters – Processes and Functions

Abiotic Components

A perspective on the abiotic processes sustaining the ecological integrity of running waters
G.E. Petts 15–27

Restoring ecological integrity of great rivers: historical hydrographs aid in defining reference conditions for the Missouri River
D.L. Galat, R. Lipkin 29–48

The river-scaling concept (RSC): a basis for ecological assessments
H.M. Habersack 49–60

Biotic Components

Biological processes in running waters and their implications for the assessment of ecological integrity
S.E. Bunn, P.M. Davies 61–70

Mayflies (*Insecta: Ephemeroptera*) and the assessment of ecological integrity: a methodological approach
E. Bauernfeind, O. Moog 71–83

Fundamentals of fish ecological integrity and their relation to the extended serial discontinuity concept
M. Jungwirth, S. Muhar, S. Schmutz 85–97

Indicators, Assessment Methods, and Applications

Assessing the ecological integrity of rivers: walking the line among ecological, political an administrative interests
O. Moog, A. Chovanec 99–109

Indicators and assessment methods for measuring the ecological integrity of semi-aquatic terrestrial environments
S.A. Innis, R.J. Naiman, S.R. Elliott 111–131

Benthos

Towards the assessment of 'ecological integrity' in running waters of the United Kingdom
D.M. Harper, J.L. Kemp, B. Vogel, M.D. Newson 133–142

The influence of environmental variables on the abundance of aquatic insects: a comparison of ordination and artificial neural networks
R. Wagner, T. Dapper, H.-H. Schmidt 143–152

Biomonitoring through biological traits of benthic macroinvertebrates: how to use species trait databases?
P. Usseglio-Polatera, M. Bournaud, P. Richoux, H. Tachet 153–162

The spatial heterogeneity of a river bottom: a key factor determining macroinvertebrate communities
J.-N. Beisel, P. Usseglio-Polatera, J.-C. Moreteau 163–171

River ecotones: carabid beetles as a tool for quality assessment
A. Boscaini, A. Franceschini, B. Maiolini 173–181

Secondary production of benthic communities at the habitat scale as a tool to assess ecological integrity in mountain streams
A. Buffagni, E. Comin 183–195

Performance of different biotic indices and sampling methods in assessing water quality in the lowland stretch of the Tiber River
A.G. Solimini, P. Gulia, M. Monfrinotti, G. Carchini 197–208

Urbanization gradients in streams of Anchorage, Alaska: a comparison of multivariate and multimetric approaches to classification
A.M. Milner, M.W. Oswood 209–223

Macroinvertebrate functional feeding group methods in ecological assessment
C. Rawer-Jost, J. Böhmer, J. Blank, H. Rahmann 225–232

The statistical power of selected indicator metrics using macroinvertebrates for assessing acidification and eutrophication of running waters
L. Sandin, R.K. Johnson 233–243

Evaluating physical habitat integrity in relation to the biological potential of streams
C.F. Rabeni 245–256

Fish

Application of an adapted index of biotic integrity to rivers of Lithuania
V. Kesminas, T. Virbickas 257–270

Fish as indicators for the assessment of the ecological integrity of large rivers
F. Schiemer 271–278

A multi-level concept for fish-based, river-type-specific assessment of ecological integrity
S. Schmutz, M. Kaufmann, B. Vogel, M. Jungwirth, S. Muhar 279–289

Variation in a Great River Index Of Biotic Integrity over a 20-year period
J.R. Gammon, T.P. Simon 291–304

Effects of residual flow and habitat fragmentation on distribution and movement of bullhead (*Cottus gobio* L.) in an alpine stream
S. Fischer, H. Kummer 305–317

Habitat condition and fish assemblage structure in a coastal mediterranean stream (Yarqon, Israel) receiving domestic effluent
S. Gafny, M. Goren, A. Gasith 319–330

Benthic invertebrate and fish communities as indicators of morphological integrity in the Enz River (south-west Germany)
W. Jansen, J. Böhmer, B. Kappus, T. Beiter, B. Breitinger, C. Hock 331–342

Abiotic Indicators

Identification of rivers with high and good habitat quality: methodological approach and applications in Austria
S. Muhar, M. Schwarz, S. Schmutz, M. Jungwirth 343–358

Using river habitat survey for environmental assessment and catchment planning in the U.K.
P.J. Raven, N.T.H Holmes, M. Naura, F.H. Dawson 359–367

Effects of catchment properties on hydrochemistry, habitat complexity and invertebrate community structure in a lowland river
B. Bis, A. Zdanowicz, M. Zalewski 369–387

Integrated Assessment Methods

Integrated ecological assessment methods as a basis for sustainable catchment management
P.F.M. Verdonschot 389–412

The development of integrated methods for assessing river conservation value
P.J. Boon 413–428

EMAP-Surface Waters: a multiassemblage, probability survey of ecological integrity in the U.S.A.
R.M. Hughes, S.G. Paulsen, J.L. Stoddard 429–443

The Austrian way of assessing the ecological integrity of running waters: a contribution to the EU Water Framework Directive
A. Chovanec, P. Jäger, M. Jungwirth, V. Koller-Kreimel, O. Moog, S. Muhar, S. Schmutz 445–452

Measuring the attainment of biological integrity in the U.S.A.: a critical element of ecological integrity
M.T. Barbour, W.F. Swietlik, S.K. Jackson, D.L. Courtemanch, S.P. Davies, C.O. Yoder 453–464

GIBSI – an integrated modelling system prototype for river basin management
A.N. Rousseau, A. Mailhot, R. Turcotte, M. Duchemin, C. Blanchette, M. Roux, N. Etong, J. Dupont, J.-P. Villeneuve 465–475

Scientific base and modular concept for comprehensive assessment of streams in Switzerland
U. Bundi, A. Peter, A. Frutiger, M. Hütte, P. Liechti, U. Sieber 477–487